

MicroBioTest Protocol

Efficacy Evaluation of Copper Enhanced Hard Surfaces as a Sanitizer - Supplemental

Testing Facility
MicroBioTest
Division of Microbac Laboratories, Inc.
105 Carpenter Drive
Sterling, VA 20164

Prepared for Luminore, Inc. 6060 Corte del Cedro Carlsbad, CA 92011

June 15, 2014

Page 17 of 33

MicroBioTest Protocol: LUM.4a.06.15.14

MicroBioTest Project: 872 . 103

TH

OBJECTIVE:

This test is designed to substantiate effectiveness claims for a substance containing copper with sanitizing claims intended to be registered with the Environmental Protection Agency as an inanimate hard surface other than those that come in contact with food or beverages. The test is consistent with the EPA Test Method for Efficacy of Copper Alloy Surfaces as a Sanitizer.

TESTING CONDITIONS:

A total of five test replicates per challenge microorganism will be evaluated using carriers prepared from the copper enhanced hard surface. Two lots of the test surface will be evaluated. Prepared carriers of the test surface will be inoculated with *Pseudomonas aeruginosa*, Methicillin Resistant *Staphylococcus aureus*, and *Escherichia coli* O157:H7 held for the stipulated contact time, transferred to a neutralizing solution and mixed. Dilutions of the neutralizer will be plated, incubated and observed for growth.

MATERIALS:

A. Test and control surfaces supplied by the sponsor: (see last page for details).

Test and control carriers: 1" x 1" coupons, also referred to as carriers

- The identity, strength, purity, and composition, or other characteristics which will appropriately define the test, control, or reference surfaces shall be determined for each batch and shall be documented by the sponsor before its use in a study. Methods of synthesis, fabrication, or derivation of the test, control, or reference surfaces shall be documented and retained by the sponsor.
- When relevant to the conduct of the study the solubility of each test, control, or reference agent shall be determined by the sponsor before the experimental start date. The stability of the test, control, or reference agent shall be determined by the sponsor before the experimental start date or concomitantly according to written standard operating procedures, which provide for periodic analysis of each batch.

Protocol: LUM.4a.06.15.14

MI

The test and control surfaces will be tested as supplied by the sponsor unless directed otherwise. All operations performed on the surfaces such as dilution or specialized storage conditions must be specified by the sponsor before initiation of testing.

The sponsor assures MicroBioTest, Division of Microbac Laboratories, Inc. (MicroBioTest) testing facility management that the test surface has been appropriately tested for identity, strength, purity, stability, and uniformity as applicable.

MicroBioTest will retain all unused test and control surfaces after completion of the test, and then only discard them with client permission in a manner that meets the approval of the safety officer.

- B. Materials supplied by MicroBioTest including but not limited to:
 - 1. Challenge microorganisms, required by EPA and the sponsor:
 - a. Pseudomonas aeruginosa, ATCC 15442
 - b. Methicillin Resistant Staphylococcus aureus (MRSA), ATCC 33592
 - c. Escherichia coli O157:H7, ATCC 35150
 - 2. Media and reagents:
 - a. Tryptic Soy Broth (TSB)
 - b. Neutralizer: 2X Letheen Broth
 - c. Phosphate Buffer Saline dilution blanks (PBS)
 - d. Tryptic Soy Agar (TSA)
 - e. Heat-inactivated Fetal Bovine Serum (FBS)
 - f. Triton X-100 solution (1% solution)
 - g. Sterile deionized water
 - h. 70-85% Isopropyl alcohol
 - 3. Miscellaneous laboratory equipment and supplies.
 - 4. Media, reagents and supplies for Antimicrobial Susceptibility Testing of MRSA:
 - a. TSA containing 5% defibrinated sheep's blood (TSA+)
 - b. 0.85% NaCl (SS)
 - c. Mueller Hinton Agar (MHA)
 - d. Control microorganism: Staphylococcus aureus, ATCC 25923

Protocol: LUM.4a.06.15.14

THE

- e. 0.5% McFarland Standard
- f. Caliper measuring device
- g. 1 μg Oxacillin disc

TEST SYSTEM IDENTIFICATION:

All test and control tube racks will be labeled with microorganism, test agent (if applicable) and project number prior to initiation of the study and during incubation. Petri dishes will be labeled with microorganism prior to initiation of the study and microorganism and project number during incubation.

EXPERIMENTAL DESIGN:

Bacteria from stock cultures will be transferred into TSB and incubated at 35-37°C for 24±2 hours. Daily transfers will be made for at least three consecutive days (but no more than 10 days). For each transfer, tubes containing 10 mL of TSB will be inoculated using two loopfuls (4-mm inside diameter) of inoculum for each tube. A 48±4 hour culture will be used for the inocula on the day of testing.

The <u>pellicle formed in the *Pseudomonas aeruginosa* culture</u> will be aspirated before use.

Transfers more than 15 days away from the stock cultures will not be used for the inocula for the test.

For each microorganism, each culture will be thoroughly mixed on a vortex-mixer and allowed to settle for ≥15 minutes. The upper two-thirds of each culture will be aspirated and used as the inoculum.

B. Addition of organic load:

To each prepared inocula, a 0.25 mL aliquot of FBS plus a 0.05 mL1% Triton X-100 solution to 4.70 mL of bacteria suspension to yield a 5% FBS and 0.01% Triton X-100 soil load.

Protocol: LUM.4a.06.15.14

C. Test and Control Carrier preparation:

The test (two lots, five replicates per lot per microorganism) and control surfaces/carriers (three replicates per microorganism) plus additional test and control surfaces as required for remaining controls (see Table 1) will be cleaned by submersion in 70-85% in Isopropyl alcohol, rinsed with sterile deionized water, and allowed to air dry. After drying completely, the carriers may be steam sterilized for 15 minutes at 121°C, or sterilized in a method approved by the sponsor. The carriers will be allowed to cool and held at ambient room temperature until use. Prior to use, each carrier will be aseptically transferred into plastic Petri dishes (one dish for each carrier) matted with two pieces of filter paper using sterile forceps.

		Test and Primary Controls			
Method	Lot	Reps	Organisms	Contact Time or Runs	TOTAL
	1	5	3	1	15
Camibinan	2	5	3	1	15
Sanitizer	3	5	3	1	15
	Control	3	3	1	9

Table 1: Test and Control Carrier Description and Count

N	/liscella	neous Cor	itrols	
NE	Viab	Sterility	TOTAL	Grand Total
3	0	1	4	19
3	0	1	4	19
3	0	1	4	19
3	3	1	7	16

D. Carrier inoculation:

A <u>0.02 mL</u> aliquot of the inoculum will be transferred onto each sterile carrier using a calibrated micropippetor. The inoculum will be spread to within approximately 1/8" of the edge of the carrier. The carriers will be allowed to dry <u>with lids ajar</u> for 30-40 minutes under ambient conditions. The exposure period (contact time) begins immediately after drying.

E. Test:

For each microorganism per lot, five inoculated and dried carriers will be held for the exposure (contact) time. The contact time will begin immediately after drying in accordance with Section D, Carrier inoculation.

At the conclusion of the contact time, each carrier will be transferred to a jar containing 20 mL of neutralizer at the appropriate staggered intervals. Each jar will be sonicated for five minutes and then rotated by hand to mix. Within one hour after sonication, serial dilutions will be prepared using PBS ($10^{-1} - 10^{-4}$). Duplicate 1.0 mL aliquots from each jar/dilution ($10^{0} - 10^{-4}$) will be plated using TSA pour

Protocol: LUM.4a.06.15.14

plates.

Plates will be incubated for 48±4 hours at 35-37°C, colonies will be counted and CFU/carrier calculated.

F. Controls:

1. Carrier quantitation control:

For each challenge microorganism, a parallel control will be run using the control carriers (surfaces) in the same manner as the test (including the contact time) with the exception that three replicates will be evaluated rather than five. All plates will be incubated appropriately in the same manner as the test plates.

2. Culture purity control:

Each prepared culture will be streaked for isolation using TSA. All plates will be incubated in the same manner as the test plates. The isolated cultures will be observed for purity.

3. Organic soil sterility control:

Duplicate 1.0 mL aliquots of the prepared organic soil will be plated in TSA pour plates. The plates will be incubated for with the test plates observed for growth or no growth.

4. Inoculum confirmation counts control:

Each prepared inoculum will be serially diluted using PBS and selected dilutions will be plated in duplicate using TSA pour plates. All plates will be incubated with the test plates.

5. Neutralizer sterility control:

A single jar of containing the neutralizer will be incubated with the test plates. The neutralizer will be observed for growth or no growth.

6. Carrier sterility control:

Protocol: LUM.4a.06.15.14

AN

An uninoculated test (per lot) and control carrier will be subcultured into independent jars containing the neutralizer and incubated with the test plates. The neutralizer will be observed for growth or no growth.

7. Carrier viability control:

For each challenge microorganism, a single inoculated <u>control carrier</u> will be subcultured into a jar containing the neutralizer and incubated with the test plates. The neutralizer jars will be observed for growth or no growth.

8. Neutralizer effectiveness control:

For each challenge microorganism, per lot of the test article, a single sterile test carrier will be neutralized in the same manner as the test (transferred into individual jars containing 20 mL of neutralizer. To each jar, a 1.0 mL aliquot of the diluted inoculum will be added to yield ≤100 CFU/mL in the neutralizer. The jar will be mixed and a 1.0 mL aliquot will be removed and plated in duplicate.

A numbers control will be performed in the same manner with the exception that a sterile control carrier will be used.

All plates will be incubated with the test plates.

9. Antimicrobial Susceptibility Testing of MRSA:

The prepared MRSA culture will be subcultured onto a TSA+ plate and the plate will be incubated for approximately 24 hours at 35-37°C. Following incubation, a suspension will be prepared by suspending growth from the TSA+ culture in SS to yield equivalent turbidity to a 0.5 McFarland Standard. This prepared suspension will be streaked onto MHA plate in a cross-hatch pattern and a 1 μ g Oxacillin disc will be placed onto the center of the plate. The plate will be inverted and incubated for \geq 24 hours at 35-37°C.

The same procedures will be conducted concurrently using the control microorganism, *Staphylococcus aureus*, ATCC 25923 to confirm the validity of the assay.

The interpretation of the zone of inhibitions (ZOI) will be based on

Protocol: LUM.4a.06.15.14

established National Committee for Clinical Laboratory Standards (NCCLS) performance standards. As currently published, (NCCLS standard M100-S21) ZOI breakpoints must be ≤ 10 mm (rounded to the nearest whole mm) confirms resistance, 11-12 mm is considered intermediate resistance, and ≥ 13 mm confirms susceptibility.

10. Microorganism confirmation procedures:

A randomly selected colony from the carrier quantitation control plates, and if applicable, a randomly selected colony from a test plate will be confirmed by colony morphology and Gram stain according to extant SOPs. The same procedures will be performed using the culture purity control plates and the result regarding purity will be documented as well.

TEST ACCEPTANCE CRITERIA:

The test will be acceptable for evaluation of the test results if the neutralizer is effective and non-toxic. The study director may consider other causes that may affect test reliability and acceptance. There are no proposed statistical methods for this test.

- The average recovery for the Carrier Quantitation Control must be at least 2.0 x 10⁴ CFU/carrier.
- The CFU recovered for the neutralizer effectiveness controls should be within 1.0 log₁₀ of the parallel neutralization confirmation control.
- The carrier sterility controls must exhibit no growth.
- The carrier viability controls must exhibit growth.
- The purity controls must demonstrate pure cultures.
- The organic soil sterility control must exhibit no growth.
- The neutralizer sterility control must exhibit no growth.
- For the Antimicrobial Susceptibility Testing: the test MRSA strain must exhibit resistance and the *Staphylococcus aureus* control strain (ATCC 25923) must exhibit susceptibility to Oxacillin.

PRODUCT EVALUATION CRITERIA:

According to EPA guidelines, the test agent meets effectiveness requirements, if the test results exhibit a bacterial reduction of at least 99.9% over the Carrier Quantitation Control.

Protocol: LUM.4a.06.15.14

Page 24 of 33

DATA PRESENTATION:

The final report will include the following information in tabular form:

- The average colony-forming units (CFU)/carrier and percent reduction for each evaluation.
- The results for all the controls.

PERSONNEL AND TESTING FACILITIES:

A study director will be assigned before initiation of the test. Resumes for technical personnel are maintained and are available on request. This study will be conducted at MicroBioTest, 105 Carpenter Drive, Sterling, VA 20164.

CONFIDENTIALITY:

All data generated at MicroBioTest are held in strictest confidence and are available only to the sponsor and the sponsor designated authorities (if applicable). In turn, no reference to MicroBioTest's promotion of the evaluated test articles may be made public by the sponsor.

REPORT FORMAT:

MicroBioTest employs a standard report format for each test design. Each final report provides the following information:

- Sponsor identification and test agent identification
- Type of test and project number
- Dates of study initiation and completion
- Interpretation of results and conclusions
- Test results
- Methods and evaluation criteria
- Signed Quality Assurance and Compliance Statements (for GLP studies, if provided by the sponsor)

Protocol: LUM.4a.06.15.14

REGULATORY COMPLIANCE AND QUALITY ASSURANCE (applicable to GLP studies only)

This study will be performed in compliance with the US Environmental Protection Agency's Good Laboratory Practices regulations, 40 CFR 160. Note: information on the identity, strength, purity, stability, uniformity, and dose solution analysis of the test agent resides with the sponsor of the study unless otherwise stated.

The Quality Assurance Unit of MicroBioTest will inspect the conduct of the study for GLP compliance. The dates of the inspections and the dates that findings are reported to the study management and study director will be included in the final report.

RECORDS TO BE MAINTAINED:

All raw data, protocol, protocol modifications, test agent records, final report, and correspondence between MicroBioTest and the sponsor will be stored in the archives at MicroBioTest, 105 Carpenter Drive, Sterling, Virginia 20164 or in a controlled facility off site.

All changes or revisions to this approved protocol will be documented, signed by the study director, dated and maintained with this protocol. The sponsor will be notified of any change, resolution, and impact on the study as soon as practical.

The proposed experimental start and termination dates; additional information about the test agent; challenge microorganism used; media and reagent identification; and the type of neutralizers employed in the test will be addressed in a project sheet issued separately for each study. The date the study director signs the protocol will be the initiation date. All project sheets will be forwarded to the study sponsor.

MAY

MISCELLANEOUS INFORMATION:

The following information is to be completed by sponsor before initiation of study:

A. Name and address:

Luminore, Inc.

6060 Corte del Cedro Carlsbad, CA 92011

B. Test surface information:

Test surface name	Luminore Copper		
Lot No.	Lot 1	Lot 2	
	441/00/	100214A	
- Manufacture Date	10/01/2014	10/02/2014	
- Expiration Date	NA	NA	
Active ingredient	Copper		
Substrate	aluminum magnesium (the sponsor will provide coupons that are not treated with the active ingredient for use as the control carriers as applicable)		

Note: both lots will be tested (therefore supplied) at or below the Lower Certified Limit (LCL)

C.	Test conditions:			
	Contact time:	(must be ≤ 2 hours))	
	Exposure temperature:	Ambient room temperature (20±1C)		
D.	Organic load – serum add	ded to achieve 5% in the inoculum:	yes	☐ no
E.	Precautions/storage - MS	SDS or certificate of analysis provided:	⊠ yes	no

Protocol: LUM.4a.06.15.14

AN

MISCELLANEOUS INFORMATION: (continued)

REPORT HANDLING: The sponsor intends to submit this information to: US EPA

Angela L. Hollingsworth

STUDY CONDUCT: GLP

PROTOCOL APPROVAL:

Sponsor:

LuminOre, Inc.

A California Corporation

Title: President

Study Director Signature:

11/2/14

MicroBioTest, A Division Observatories Inc. 105 Carpenter or Sterling, Virginia 20164

Date Issued: 11/03/14 P	roject Sheet No. 1 Page		ry Project Identification I		
STUDY TITLE: Efficacy Evaluation of Copper		STUDY DIRECTOR: Angela L. Hollingsworth			
Enhanced Hard Surfaces as a Sanitizer - Supplemental		ag Joels 1/3/14			
		Signature /		Date	
TEST AND CONTROL ART	ICLES:	LOT NO:	DATE RECEIVED:	DS NO:	
LuminOre Copper		100214A	10/17/14	E526	
LuminOre Copper		100114A	10/17/14	E527	
Non Treated Coupons		Not applicable	10/17/14	E532 -	
PERFORMING DEPARTMENT:			STORAGE CONDITIONS: Location: K4		
Applied Microbiology Laboratory			■ Dark ■ Ambient Room Temperature		
			reezer □ Refrigerator □		
PROTECTIVE PRECAUTIO			of Analysis was provide	ed)	
PHYSICAL DESCRIPTION:					
PURPOSE: See attached p					
PROPOSED EXPERIMENT			DATE: 11/06/14		
CONDUCT OF STUDY:			N: Thomas J. Val	onto	
SPONSOR: Luminore, Inc		CONTACT PERSOI Phone:	(760) 431-770		
6060 Corte d		E-mail:	tom@luminore		
Carlsbad, CA	92011	E-IIIaII.	tornwiaminore	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
TEST CONDITIONS:		1			
Challenge organism(s):	Pseudomonas aeruginosa Methicillin Resistant Staph Escherichia coli O157:H7,	ylococcus aureus, AT	CC 33592		
Active ingredient(s):	Copper				
Neutralizer(s):	Letheen Broth - 2X				
Contact Time(s):	2 hours				
Contact Temperature(s):	Ambient (20±1°C)	Dilution(s)	Ready to Use		
Organic Load:	■ Yes / □ No (Per the pro	tocol to achieve 5% in	n the inoculum)		
Incubation Time(s):	48±4 hours				
Incubation Temperature(s):	35-37°C				
Comments: The sponsor provided alum active ingredient for use as					

Micro	oBioTest, A Division of incrobac Laboratories II	nc. Too Carpenter of.	Sterling, Virginia 20164	
Date Issued:	11/03/14 Project Sheet No. 2 Page	No. 1 Laborator	y Project Identification	No. 872-103
STUDY TITL	E: Efficacy Evaluation of Copper	STUDY DIRECTO	R: Angela L. Hollingsw	orth
	ard Surfaces as a Sanitizer - Supplemental	011	0001	In L.
		- 4	() () () () ()	13/19
		Signature		Date
TEST AND C	ONTROL ARTICLES:	LOT NO:	DATE RECEIVED:	DS NO:
LuminOre Co	pper	100214A	10/17/14	E526
LuminOre Copper		100114A	10/17/14	E527
Non Treated	Coupons	Not applicable 10/17/14 E532		
The state of the s			TIONS: Location: K4	
Applied Micro	bbiology Laboratory		Room Temperature	
			reezer □ Refrigerator [1 Other:
	PF STUDY: □ FDA ■ EPA □ R&D ■ GLP I			
SPONSOR:	Luminore, Inc.	CONTACT PERSON		
	6060 Corte del Cedro	Phone:	(760) 431-770	
	Carlsbad, CA 92011	E-mail:	tom@luminore	e.com
EXPLANATION	ON:			
	te the units of measurement for the cordment serves to clarify an omission in the p		act time will be 2 hot	urs. This

MicroBioTest, A Division of Microbac Laboratories Inc. 105 Carpenter Di. Sterling, Virginia 20164

Date Issued: 02/04/15 Project Sheet No. 3 Page	No 1 Laborate	ory Project Identification	No. 872-103
STUDY TITLE: Efficacy Evaluation of Copper		OR: Angela L. Hollingsw	
Enhanced Hard Surfaces as a Sanitizer - Supplemental	J. B	Defrace FEB	0 4 2 015 Date
TEST AND CONTROL ARTICLES:	LOT NO:	DATE RECEIVED:	DS NO:
LuminOre Copper	100214A	10/17/14	E526
LuminOre Copper	100214A	10/17/14	E527
Non Treated Coupons	Not applicable	10/17/14	E532
PERFORMING DEPARTMENT:		DITIONS: Location: K4	L002
pplied Microbiology Laboratory ■ Dark ■ Ambient Room Temperature □ Desiccator □ Freezer □ Refrigerator □			□ Other:
CONDUCT OF STUDY: ☐ FDA ■ EPA ☐ R&D ■ GLP ☐	☐ GCP ☐ Other:		
SPONSOR: Luminore, Inc.	CONTACT PERSO		
6060 Corte del Cedro	Phone:	(760) 431-770	
Carlsbad, CA 92011	E-mail:	tom@luminore	e.com
Protocol Amendment(s): 2. Page 11 of the protocol did not specify the perpercentage of copper in each test surface is a • Lot No. 100214A: 81.9% copper • Lot No. 100114A: 81.0% copper		ach test surface. Per th	e sponsor, the



6060 Corte Del Cedro Carlsbad, CA 92011 Phone: 760-431-7705

Fax: 760-431-7706

Certificate of Analysis

Material: Copper Batch No.: 100214A

Parameter	Actual	Acceptable
Temperature	73°F	No less than 70°F
Humidity	33%	Less than or equal to 40%
Cross link	Pass	Pass
Mil thickness	Pass	No Less than 12 mils

Passes all quality control standards for the LuminOre product.

APPROVED BY:

Date: 10-16-14



6060 Corte Del Cedro Carlsbad, CA 92011 Phone: 760-431-7705

Fax: 760-431-7706

Date: 10-16-14

Certificate of Analysis

Material: Copper Batch No.: 100114A

Parameter	Actual	Acceptable
Temperature	79°F	No less than 70°F
Humidity	36%	Less than or equal to 40%
Cross link	Pass	Pass
Mil thickness	Pass	No Less than 12 mils

Passes all quality control standards for the LuminOre product.

APPROVED BY: